

Accessing pion GPDs through the Sullivan process: is it feasible?

J. M. Morgado

Dpto. Ciencias Integradas and CEAFCM, Universidad de Huelva, E-21071 Huelva, Spain.

V. Bertone

Irfu, CEA, Université Paris-Saclay, 91191, Gif-sur-Yvette, France.

M. Defurne

Irfu, CEA, Université Paris-Saclay, 91191, Gif-sur-Yvette, France.

F. De Soto

Dpto. Sistemas Físicos, Químicos y Naturales, Universidad Pablo de Olavide, E-41013 Sevilla, Spain.

C. Mezrag

Irfu, CEA, Université Paris-Saclay, 91191, Gif-sur-Yvette, France.

H. Moutarde

Irfu, CEA, Université Paris-Saclay, 91191, Gif-sur-Yvette, France.

J. Rodríguez-Quintero

Dpto. Ciencias Integradas and CEAFCM, Universidad de Huelva, E-21071 Huelva, Spain.

J. Segovia

Dpto. Sistemas Físicos, Químicos y Naturales, Universidad Pablo de Olavide, E-41013 Sevilla, Spain.

Received 14 January 2022; accepted 25 February 2022

Describing hadronic structure is one of the most intriguing problems in physics. In this respect, generalized parton distributions (GPDs) constitute an outstanding tool, allowing to draw “three dimensional pictures” of hadron’s inside. Starting from contemporary models for pion’s GPDs fulfilling all constraints imposed by QCD, we compute Compton form factors of pions subjected to deeply virtual Compton scattering. We show CFF’s behaviour to be gluon-dominated at EIC’s kinematics. Finally we evaluate lepton-beam-spin asymmetries in the Sullivan process, demonstrating the existence of such and thus triggering optimism about the possibility of probing pion’s 3D structure at electron-ion colliders.

Keywords: Non-perturbative QCD; DVCS; CFFs; EIC; sullivan process.

DOI: <https://doi.org/10.31349/SuplRevMexFis.3.0308099>

1. Introduction

Among all hadrons, pions are one of the most intriguing systems of study [1,2]. As Nambu-Goldstone bosons of QCD’s dynamical breakdown of chiral symmetry, they constitute a unique laboratory for the study of emergent strong phenomena. Moreover, as “two-bodies” bound-states they are the simplest systems tied by the strong interaction. Therefore, gaining insights into pion’s structure is of great interest [1,3-7] and the most direct tool we have for this purpose is lepton scattering.

Unfortunately targeting pions is a really challenging task, requiring indirect approaches to be explored in order to probe pion’s structure. In this respect, the so-called Sullivan process [8] provides a remarkable tool. In a nutshell, given that the necessary conditions are met (see Sec. 2), the paradigmatic case of electron-proton scattering can also take place through Deeply Virtual Compton Scattering (DVCS) with a

pion in the meson cloud of the nucleon; opening a clear window to the study of pion’s structure. Moreover two mechanisms contribute to such process [9]: Bethe-Heitler (BH) and DVCS, which is related with Generalized Parton Distributions (GPDs) [10].

GPDs were introduced more than two decades ago as extensions for the “classical” Parton Distribution Functions (PDFs) to off-forward kinematics [10,13]. They were shown to offer access to the three-dimensional structure of hadrons [14] and a direct connection with QCD’s fundamental degrees of freedom through the energy-momentum tensor [15].

As interesting as they are, GPDs are notoriously difficult to extract from experiment and theory [16,17]. The first models for pion GPDs fulfilling all the requirements from quantum field theory are contemporary [18,19], but the possibility of accessing them at future experimental facilities demands further scrutiny. We herein report on a recent effort addressing this last point [20].

2. Accessing pion GPDs: the Sullivan process

As shown in a seminal paper by J. D. Sullivan [8], the cross-section for exclusive electron-nucleon scattering in the Bjorken limit receives sizeable contributions from pion-nucleon final states when the momentum transferred between the initial and final nucleon states, t , remains small and near the threshold for pion production: $|t| \lesssim 0.6 \text{ GeV}^2$ [21]. Under these conditions, such process can be interpreted in the so-called *one-pion-exchange* approximation, *i.e.* in terms of pion-photon scattering amplitudes. Moreover, the invariant mass of the nucleon- π system being cut as $M_{N\pi}^2 \gtrsim 4 \text{ GeV}^2$, possible contamination from nucleon resonances is minimized [8,9], allowing to interpret the Sullivan process in terms of direct interaction between a photon and a slightly off-shell pion.

Following these arguments, the amplitude for the Sullivan process ($ep \rightarrow e\pi n\gamma$) can be “factorized” as:

$$\mathcal{M}_{ep \rightarrow e\pi n\gamma} = \mathcal{M}_{p \rightarrow \pi n} \mathcal{M}_{e\pi \rightarrow e\pi\gamma}, \quad (1)$$

drawing a clear path toward the experimental analysis of pion’s structure through the amplitude for $e\pi \rightarrow e\pi\gamma$.

Two subprocesses give contributions to such amplitude [9]: Bethe-Heitler scattering, which provides direct access to the pion’s Electromagnetic Form Factor (EFF) and has already been exploited for its extraction at large photon virtualities [22]. And a DVCS contribution:

$$\mathcal{M}_{e\pi \rightarrow e\pi\gamma} = \mathcal{M}_{\text{BH}} + \mathcal{M}_{\text{DVCS}}. \quad (2)$$

DVCS amplitudes are parametrized by Compton Form Factors (CFFs) [10]. These being defined as convolutions of a perturbatively calculable kernel with GPDs [12,23-26], the DVCS contribution to $e\pi \rightarrow e\pi\gamma$ allows for a direct study of pion’s three-dimensional structure.

Cross sections being defined as the modulus squared of the corresponding amplitude, here Eq. (2), three contributions to that of the Sullivan process are identified: pure DVCS (BH), parametrized by modulus squared CFFs (EFFs); and a third contribution, the interference term between two such subprocesses, directly related with the real and imaginary parts of the CFFs [9,23]. Moreover, at leading twist, pure DVCS and BH contributions exhibit no dependence on lepton-beam-spin polarization while the interference term does [9]. Therefore, the lepton’s Beam Spin Asymmetry (BSA), \mathcal{A} :

$$\mathcal{A} := \frac{\sigma^\uparrow - \sigma^\downarrow}{\sigma^\uparrow + \sigma^\downarrow} \Big|_{e\pi \rightarrow e\pi\gamma} = \frac{\mathcal{S}_{\text{Im}} \text{Im}(\mathcal{H}_\pi)}{\text{BH} + \text{DVCS} + \mathcal{C}_{\text{Re}} \text{Re}(\mathcal{H}_\pi)}, \quad (3)$$

grants access to the real and imaginary parts of CFFs, \mathcal{H}_π ; thus allowing to employ the Sullivan process to measure them (at least formally) or, equivalently, to gain insights into the pion’s three-dimensional structure through GPDs.

Furthermore, the coefficients of each term in Eq. (3) are shown to be proportional to $\cos \phi_\pi$ (\mathcal{C}_{Re}) and $\sin \phi_\pi$ (\mathcal{S}_{Re})

[9,27], with ϕ_π the angle between the planes defined by (incoming and outgoing) leptons and pions in the center of mass frame of $\pi\gamma$ in the final state [28]. This means that not only ratios $\text{Im}(\mathcal{H}_\pi)/\text{Re}(\mathcal{H}_\pi)$ can be measured through the Sullivan process, but also to exploit the ϕ_π dependence of its interference to separately access the real and imaginary parts of CFFs.

The clarity of the above described procedure does not preclude the feasibility of a practical analysis of pion’s structure to be obscured by experimental artefacts. Given that various experimental facilities devoted to the study of hadron structure are planned [5-7], a feasibility study of the methodology presented herein is timely.

Thus we choose to perform an analysis of the Sullivan process at the future Electron-Ion Collider (EIC) [5]. Relying on state-of-the-art models for pion GPDs [19], we evaluate Compton form factors at Next-to-Leading Order (NLO) in the strong coupling constant. From their analysis we demonstrate that pion’s response to Compton scattering in the Bjorken limit is gluon-dominated. Finally we compute lepton-beam-spin asymmetries, showing that a clearly non-zero asymmetry might be expected at the foreseen EIC, thus triggering optimism about the possibility of probing pion’s three-dimensional structure [20].

3. Modelling of pion GPDs

The first ingredient needed for the present study is a reliable model for pion’s 3D structure. However, modelling of hadron GPDs is a difficult task. Mainly because their dependence on: x , average momentum fraction of the active quark; ξ , half the longitudinal momentum transferred; and t_π , squared momentum transfer between pion states, are constrained by a set of properties [13,29]. In fact, only modern strategies have been able to build models fulfilling all of them by construction [18,19]. It is precisely such approach to which we stick.

In summary, under the assumption of chiral symmetry, DGLAP GPDs ($|x| \geq |\xi|$) can be built from the usual PDFs [19,30,31]; with the positivity property granted [19]. Then, the covariant extension formalism [18,32] can be employed for the kinematic completion of the model within the ERBL domain ($|x| \leq |\xi|$); polynomiality and support properties being also satisfied (Fig. 1).

According to [19], modelling of pion GPDs is therefore reduced (in a first stage) to that of the PDF. In this respect we choose to start from the results presented in Ref. [33], where realistic *Ansätze* for the pion’s PDF were obtained from continuum Schwinger methods, showing a large- x behaviour which remains compatible with that extracted from experimental data including soft-gluon (threshold) resummation [34-36].

The corresponding DGLAP GPD model was presented in Ref. [31], and further elaborated with its kinematic completion in Ref. [19]. It was also shown to fulfil by construction with the support property of GPDs [37]; with the bounds




FIGURE 1. Diagram describing contemporary modelling strategies for pion GPDs allowing to fulfil every fundamental property required by quantum field theory as introduced in Ref. [19].

imposed by positivity of the underlying Hilbert-space norm [38-41], and also to preserve Lorentz invariance, as exhibited through polynomiality [42-44], they all general properties of GPDs. Furthermore, they have been also shown to account for partial conservation of the axial current [45,46], as one might expect in the case of the pion. Moreover, the resulting model exhibits agreement with available extractions for electromagnetic [22,47] and Gravitational Form Factors (GFFs) [48] of the pion, specially in the low- t_π domain. Even more, the GPD thus obtained showed to be continuous along the $x = \xi$ line, as required by analyticity, for the amplitudes of DVCS to be finite [49,50].

The resulting model puts the cap on a long effort developed during the last decade [18,32,46,51-53], showing that a realistic description of pion's 3D structure can be achieved from continuum Dyson-Schwinger computations. The model presented in Ref. [19] exhibiting not only the formal requirements needed on a GPD, but also those of a phenomenological origin, constitutes a serious candidate for a realistic description of pion's structure. Its foundations on very first principles of quantum field theory together with its agreement with existing data for observables like EFFs, GFFs (specially in the region to be probed at EICs) or PDFs, points toward the possibility of obtaining solid predictions on the phenomenology to be observed at future colliders. Its use in the present study being self-justified and confidence on the model's independence of the results pushed forward.

3.1. GPD evolution

Given a formal description of pion's structure through GPDs, facing an analysis of the phenomenology to be observed at the foreseen EIC crucially requires QCD's scale-evolution to be handled. The EIC will operate at energies in the range of tens of GeV^2 s [5], therefore running the GPD model from its definition scale to an experimentally relevant one is mandatory.

As described by [19,31], in the sense of scale-dependence, the model was built under a sole assumption:

the existence of a scale, μ_H , at which the entire parton content of the pion is described by two dressed valence-quarks. Such hypothesis has proved to yield successful descriptions at the PDF level [33,54]. Moreover, in the case of the pion, it has been recently found that its existence is a solid prediction of available Lattice-QCD and Drell-Yan data [31,36].

For these reasons we choose a description of pion's structure to be given at a reference scale by the u - and \bar{d} -GPDs developed in Ref. [19]. We take GPD scale-evolution to be driven by the process independent effective coupling introduced in Ref. [55], which accounts for saturation in the infrared regime by means of dynamical generation for a gluon mass-scale. Thus, the reference scale is set accordingly to $\mu_H = 0.331 \text{ GeV}$ [54], and the Leading Order (LO) QCD evolutions for GPDs solved through `Apfel++` software [56,57].

4. Compton form factors

With a realistic model for pion GPDs evolved to experimentally relevant energy scales, we are now in a position to em-




FIGURE 2. DVCS Compton Form Factors at a factorization scale of 2 GeV^2 . *Upper panel* - Real part. *Lower panel* - Imaginary part. *Legend* - Red line: LO evaluation; blue line: NLO without gluon GPDs. Black line: full NLO result.




FIGURE 3. Expected lepton-beam-spin asymmetries as a function of Q^2 for $\xi \in [10^{-3}, 10^{-2}]$. Legend - Red circles: LO evaluation; blue triangles: NLO without the gluon GPDs. Black squares: full NLO result. (data taken from [20]).

ploy them in computing CFFs, \mathcal{H}_π , the coefficient functions parametrizing the amplitudes for DVCS [10,24,58,59]:

$$\mathcal{H}_\pi(\xi, t, Q^2) = \sum_{i=\{q\},g} \int_{-1}^1 \frac{dx}{\xi} \mathcal{K}^i\left(\frac{x}{\xi}, \frac{Q^2}{\mu_F^2}\right) \times H_\pi^i(x, \xi, t_\pi; \mu_F^2), \quad (4)$$

with the sum extended over all possible quark flavours and gluons; and where \mathcal{K}^i is a convolution kernel calculable in perturbation theory.

Interestingly, the gluon-type contribution to the convolution kernel, $\mathcal{K}^g(z)$, vanishes at LO. It is only from NLO on contributions in the strong running coupling when gluon content within the probed hadron enters the game [24,59-61], affecting the behaviour of the hadrons subjected to deeply virtual Compton scattering. For this reason we compute the pion's DVCS CFFs through three different approaches: (i) at LO, (ii) at NLO and finally, (iii) at NLO but manually setting the gluon contribution to zero.

With the help of PARTONS framework [62], calculation of CFFs through these three approaches is feasible. As an illustration, results at a factorization scale of 2 GeV^2 are shown in Fig. 2. There, it is clearly seen that both LO and NLO without gluon calculations, yield very similar results. The real and imaginary parts of the CFFs being positive definite and decreasing functions of the skewness. On the contrary, the full NLO result shows that gluon content within the pion gives the dominant contribution to the DVCS CFF. Specially in the low- ξ region, which is the one to be probed at the EIC [5]. Crucially, the observed behaviour drastically changes. Even flipping sign for both, real and imaginary parts. In this respect, it is important to notice that gluon dominance yield a CFF roughly behaving as $\xi^{-1.4}$. A result which remains compatible with those independently obtained from DVCS dispersion relations with one subtraction constant [63]; once again supporting, as argued in Sec. 3, supporting a limited model dependence of our predictions.

5. Beam-spin asymmetries at the EIC

Finally, after the development of the preceding sections, we are in a position to evaluate the cross-section for the Sulli-

van process. To this end we build a Monte Carlo event generation algorithm. We account for the prescriptions given in Ref. [9], in charge of assuring the interpretation of the process in the one-pion-exchange approximation; and also the EIC's detector characteristics specified in Ref. [5]. In that way, the phase-space of kinematic configurations is generated, and the formulae given in Ref. [9] exploited for the corresponding evaluation of the cross-sectionsⁱ. Finally, lepton's BSA are computed as defined in Eq. (3).

The obtained results are shown in Fig. 3 for $\xi \in [10^{-3}, 10^{-2}]$ and four different bins in Q^2 . In agreement with those results discussed in Sec. 4, LO and NLO without gluon contribution asymmetries are nearly indistinguishable. As expected, a sinusoidal shape is observed for the asymmetry at all explored scales [29]. Only an enhancement in their amplitudes is observed as the virtuality of the probing photon increases. In contrast, the result yielded by the full NLO calculation exhibits a markedly different behaviour. At low Q^2 , as a manifestation of gluon-dominance, the asymmetry changes sign with respect to the LO and NLO/NoGluon ones. As photon's virtuality increases, its amplitude is now reduced; turning compatible with zero at moderate scales. Moreover, at high enough Q^2 , the NLO result approaches the tendency of LO results, as expected from perturbation theory.

6. Conclusions

In the light of the results discussed herein, two main conclusions can be drawn. First, gluon content within the pion gives the dominant contribution to its behaviour in DVCS at low virtuality and small ξ . Second, a clearly non-vanishing lepton-beam-spin asymmetry is to be observed at the future EIC, thus opening the possibility to experimentally access, for the first time, pion's three dimensional structure; and even allowing to pin down the regime of gluon-dominance through an observable sign change.

Acknowledgements

We would like to thank P. Barry, H. Dutrieux, T. Meisgny, B. Pire, K. Raya, C.D. Roberts, D. Sokhan, Q.-T. Song, P. Sznajder and J. Wagner for interesting discussions and stimulating

comments. J.M.M.C acknowledges support from University of Huelva (grant EPIT-2021). F.S., J.R.Q. and J.S.'s work is supported by Ministerio de Ciencia e Innovación (Spain) under grant PID2019-107844GB-C22; Junta de Andalucía, under contract No. operativo FEDER Andalucía 2014-2020UHU-1264517 and P18-FR-5057 and PAIDI FQM-370.

This work was also supported by the European Union's Horizon 2020 research and innovation programme under grant agreement No 824093; and through the GLUODYNAMICS project funded by "P2IO LabEx (ANR-10-LABX-0038)" in the framework "Investissements d'Avenir" managed by the Agence Nationale de la Recherche (ANR), France.

- i.* For further details see [20].
1. A. C. Aguilar *et al.*, Pion and Kaon Structure at the Electron-Ion Collider. *Eur. Phys. J. A*, **55** (2019)190.
 2. C. D. Roberts, D. G. Richards, T. Horn, and L. Chang, Insights into the emergence of mass from studies of pion and kaon structure. *Prog. Part. Nucl. Phys.*, **120** (2021) 103883.
 3. R. A. Montgomery *et al.*, Proposed measurement of tagged deep inelastic scattering in Hall A of Jefferson lab. *AIP Conf. Proc.*, **1819** (2017) 030004.
 4. J. Arrington *et al.*, Revealing the structure of light pseudoscalar mesons at the electron-ion collider. *J. Phys. G*, **48** (2021) 075106.
 5. R. Abdul-Khalek *et al.*, Science Requirements and Detector Concepts for the Electron-Ion Collider: EIC Yellow Report. arxiv:2103.05419, 3 (2021).
 6. D. P. Anderle *et al.*, Electron-ion collider in China. *Front. Phys. (Beijing)*, **16** (2021) 64701.
 7. B. Adams *et al.*, Letter of Intent: A New QCD facility at the M2 beam line of the CERN SPS (COMPASS+/AMBER). **8** (2018).
 8. J. D. Sullivan. One pion exchange and deep inelastic electron-nucleon scattering. *Phys. Rev. D*, **5** (1972) 1732.
 9. D. Amrath, M. Diehl, and J.-P. Lansberg, Deeply virtual Compton scattering on a virtual pion target. *Eur. Phys. J., C* **58** (2008) 179.
 10. X. Ji, Deeply virtual Compton scattering. *Phys. Rev. D* **55** (1997) 7114.
 11. D. Müeller, D. Robaschik, B. Geyer, F. M. Dittes, and J. Horejsi, Wave functions, evolution equations and evolution kernels from light ray operators of QCD. *Fortsch. Phys.* **42** (1994) 101.
 12. A.V. Radyushkin. Nonforward parton distributions. *Phys. Rev. D* **56** (1997) 5524.
 13. M. Diehl. Generalized parton distributions. *Phys. Rept.* **388** (2003) 41.
 14. M. Burkardt. Impact parameter dependent parton distributions and off forward parton distributions for. *Phys. Rev. D* **62** (2000) 071503. [Erratum: *Phys. Rev.D*66,119903(2002)].
 15. X. Ji. Gauge-Invariant Decomposition of Nucleon Spin. *Phys. Rev. Lett.*, **78** (1997) 610.
 16. V. Bertone, H. Dutrieux, C. Mezrag, H. Moutarde, and P. Sznajder, Deconvolution problem of deeply virtual Compton scattering. *Phys. Rev. D*, **103** (2021) 114019.
 17. V. Bertone, H. Dutrieux, C. Mezrag, H. Moutarde, and P. Sznajder, Shadow generalized parton distributions: a practical approach to the deconvolution problem of DVCS. In 28th International Workshop on Deep Inelastic Scattering and Related Subjects, 7 (2021).
 18. N. Chouika, C. Mezrag, H. Moutarde, and J. Rodríguez-Quintero, Covariant Extension of the GPD overlap representation at low Fock states. *Eur. Phys. J.C* **77** (2017) 906.
 19. J. M. Morgado *et al.*, Pion generalized parton distributions: A path toward phenomenology, *Phys. Rev. D* **105** (2022) 094012. <https://doi.org/10.1103/PhysRevD.105.094012>
 20. J. M. Morgado *et al.*, Accessing the pion 3D structure at US and China Electron-Ion Colliders. *Phys. Rev. Lett.* **128** (2022) 202501. <https://doi.org/10.1103/PhysRevLett.128.202501>.
 21. S.-X. Qin, C. Chen, C. Mezrag, and C. D. Roberts, Offshell persistence of composite pions and kaons. *Phys. Rev., C* **97** (2018) 015203.
 22. G. M. Huber *et al.*, Charged pion form-factor between $Q^2 = 0.60\text{-GeV}^2$ and 2.45-GeV^2 . II. Determination of, and results for, the pion form-factor. *Phys. Rev. C* **78** (2008) 045203.
 23. A. V. Belitsky and D. Müeller, Exclusive electroproduction revisited: treating kinematical effects. *Phys. Rev. D* **82** (2010) 074010.
 24. B. Pire, L. Szymanowski, and J. Wagner, NLO corrections to timelike, spacelike and double deeply virtual Compton scattering. *Phys.Rev. D* **83** (2011) 034009.
 25. J. C. Collins and A. Freund, Proof of factorization for deeply virtual Compton scattering in QCD. *Phys. Rev. D* **59** (1999) 074009.
 26. X. Ji and J. Osborne, One loop corrections and all order factorization in deeply virtual Compton scattering. *Phys. Rev. D* **58** (1998) 094018.
 27. A. V. Belitsky, D. Müeller, and A. Kirchner, Theory of deeply virtual Compton scattering on the nucleon. *Nucl. Phys. B* **629** (2002) 323.
 28. A. Bacchetta, U. D'Alesio, M. Diehl, and C. A. Miller. Single-spin asymmetries: The Trento conventions. *Phys. Rev. D* **7** (2004) 117504.
 29. A.V. Belitsky and A.V. Radyushkin. Unraveling hadron structure with generalized parton distributions. *Phys. Rept.* **418** (2005) 1.
 30. J.-L. Zhang *et al.*, Measures of pion and kaon structure from generalised parton distributions. *Phys. Lett. B*, **815** (2021) 136158.

31. K. Raya, Z.-F. Cui, L. Chang, J. M. Morgado, C. D. Roberts, and J. Rodríguez-Quintero. Revealing pion and kaon structure via generalised parton distributions. *arXiv:2109.11686*, 9 (2021).
32. N. Chouika, C. Mezrag, H. Moutarde, and J. Rodríguez-Quintero. A Nakanishi-based model illustrating the covariant extension of the pion GPD overlap representation and its ambiguities. *Phys. Lett. B* **780** (2018) 287.
33. M. Ding, K. Raya, D. Binosi, L. Chang, C. D. Roberts, and S. M. Schmidt, Symmetry, symmetry breaking, and pion parton distributions. *Phys. Rev. D*, **101** (2020) 054014.
34. P. C. Barry, Chueng-Ryong Ji, N. Sato, and W. Melnitchouk. Global QCD analysis of pion parton distributions with threshold resummation. *arXiv:2108.05822*, **8** (2021).
35. M. Aicher, A. Schafer, and W. Vogelsang. Soft-gluon resummation and the valence parton distribution function of the pion. *Phys. Rev. Lett.*, **105** (2010) 252003.
36. Z. F. Cui *et al.*, *Concerning pion parton distributions*. **12** (2021).
37. M. Diehl and T. Gousset., Time ordering in off diagonal parton distributions. *Phys. Lett., B* **428** (1998) 359.
38. B. Pire, J. Soffer, and O. Teryaev, Positivity constraints for off-forward parton distributions. *Eur. Phys. J. C* **8** (1999) 103.
39. A.V. Radyushkin, Double distributions and evolution equations. *Phys. Rev. D* **59** (1999) 014030.
40. M. Diehl, T. Feldmann, R. Jakob, and P. Kroll, The Overlap representation of skewed quark and gluon distributions. *Nucl. Phys. B* **596** (2001) 33.
41. P. V. Pobylitsa, Positivity bounds on generalized parton distributions in impact parameter representation. *Phys. Rev. D*, **66** (2002) 094002.
42. X. Ji, Off forward parton distributions. *J. Phys. G* **24** (1998) 1181.
43. A.V. Radyushkin. Symmetries and structure of skewed and double distributions. *Phys. Lett. B* **449** (1999) 81.
44. M. V. Polyakov and C. Weiss. Skewed and double distributions in pion and nucleon. *Phys. Rev. D* **60** (1999) 114017.
45. M. V. Polyakov. Hard exclusive electroproduction of two pions and their resonances. *Nucl. Phys. B* **555** (1999) 231.
46. C. Mezrag *et al.*, Sketching the pion's valence-quark generalised parton distribution. *Phys. Lett. B* **741** (2014)190.
47. S.R. Amendolia *et al.*, A Measurement of the Space - Like Pion Electromagnetic Form-Factor. *Nucl. Phys. B* **277** (1986) 168.
48. S. Kumano, Q.-T. Song, and O. Teryaev, Hadron tomography by generalized distribution amplitudes in pion-pair production process and gravitational form factors for pion. *Phys. Rev. D* **97** (2018) 014020.
49. J. Collins. The non-triviality of the vacuum in light-front quantization: An elementary treatment. *arXiv:1801.03960*, 1 (2018).
50. V. Bertone, H. Dutrieux, C. Mezrag, J. M. Morgado Chávez, and H. Moutarde, Revisiting evolution equations for generalised parton distributions (2022).
51. C. Mezrag, H. Moutarde, and F. Sabati'e. Test of two new parameterizations of the Generalized Parton Distribution H. *Phys. Rev. D* **88** (2013) 014001.
52. C. Mezrag, H. Moutarde, J. Rodríguez-Quintero, and F. Sabatiè. Towards a Pion Generalized Parton Distribution Model from Dyson-Schwinger Equations. *arXiv:1406.7425*, (2014).
53. C. Mezrag, H. Moutarde, and J. Rodríguez-Quintero. From Bethe-SalpeterWave functions to Generalised Parton Distributions. *Few Body Syst.* **57** (2016) 729.
54. Z.-F. Cui *et al.*, Kaon and pion parton distributions. *Eur. Phys. J. C*, **80** (2020) 1064.
55. J. Rodríguez-Quintero, D. Binosi, C. Mezrag, J. Papavassiliou, and C. D. Roberts, Process-independent effective coupling. From QCD Green's functions to phenomenology. *Few Body Syst.*, **59** (2018) 121.
56. V. Bertone, S. Carrazza, and J. Rojo, APFEL: A PDF Evolution Library with QED corrections. *Comput. Phys. Commun.*, **185** (2014) 1647.
57. V. Bertone, APFEL++: A new PDF evolution library in C++. *PoS, DIS2017:201*, (2018).
58. K. Kumericki, S. Liuti, and H. Moutarde, GPD phenomenology and DVCS fitting. *Eur. Phys. J., A* **52** (2016) 157.
59. H. Moutarde, B. Pire, F. Sabatie, L. Szymanowski, and J. Wagner, On timelike and spacelike deeply virtual Compton scattering at next to leading order. *Phys. Rev. D* **87** (2013) 054029.
60. X. Ji and J. Osborne, One loop QCD corrections to deeply virtual Compton scattering: The Parton helicity independent case. *Phys. Rev. D* **57** (1998) 1337.
61. A. V. Belitsky, D. Mueller, L. Niedermeier, and A. Schafer, Deeply virtual Compton scattering in next-to-leading order. *Phys. Lett. B* **474** (2000) 163.
62. B. Berthou, PARTONS: PARTonic Tomography Of Nucleon Software. A computing framework for the phenomenology of Generalized Parton Distributions. *Eur. Phys. J. C* **78** (2018) 478.
63. M. Diehl and D. Yu. Ivanov, Dispersion representations for hard exclusive processes: beyond the Born approximation. *Eur. Phys. J. C*, **52** (2007) 919.