

Bright soliton of Stochastic perturbed Biswas-Milovic equation with cubic-quintic-septic law having multiplicative white noise

N. Ozdemir^a, S. Altun^b, M. Ozisik^c, A. Secer^d, and M. Bayram^e,

^a*Software Engineering, Istanbul Gelisim University, Istanbul-Turkey,*
e-mail: neozemir@gelisim.edu.tr

^b*Suleymaniye Mh. Destan Sk., Bursa-Turkey,*
e-mail: selvialtun89@gmail.com

^c*Mathematical Engineering, Yildiz Technical University, Istanbul-Turkey,*
e-mail: ozisik@yildiz.edu.tr

^d*Computer Engineering, Biruni University, Istanbul-Turkey,*
e-mail: asecer@biruni.edu.tr; mustafabayram@biruni.edu.tr

Received 15 August 2023; accepted 23 October 2023

For the first time, the adopted stochastic form of the perturbed Biswas-Milovic equation with cubic-quintic-septic law having spatio-temporal and chromatic dispersion in the presence of multiplicative white noise in Itô sense was presented and examined. The Biswas-Milovic equation models numerous physical phenomena occurring in optical fiber. We analyzed the optical soliton solutions of the stochastic model with the aid of a subversion of the new extended auxiliary equation method. Furthermore, we investigated the evaluation of the noise impacts and the effects of some model parameters on the dynamics of the generated soliton. Finally, graphical depictions of the derived soliton types were represented for some solution functions. The stochastic model and the derived results will contribute to the comprehension of the nonlinear dynamics of pulse propagation in optical fibers which has great importance for the advancement of optical communication engineering.

Keywords: Stochastic model; Wiener process; Noise strength; Brownian motion.

DOI: <https://doi.org/10.31349/RevMexFis.70.021303>

1. Introduction

The nonlinear stochastic partial differential equations (SPDEs) consist of stochastic derivatives or noise terms. The noise terms represent the random fluctuations or uncertainties inherent in the system being modeled. SPDEs are of great importance in the modeling of events with optical fibers, especially in optical communication. The existence of stochastic terms gives rise to extra challenges and needs specialized approaches for their analysis and solution. So, interpreting physically the SPDEs, optical soliton solutions of SPDEs have lately become the focus of attention in the literature. In particular, adopted stochastic models are constantly being developed in the literature in terms of models developed in the fields of telecommunication, fibers, nano-fibers, optics, and optoelectronics due to the importance of optical solitons.

The nonlinear Schrödinger equations (NLSEs) have a principal role in engineering and various fields [1]. Therefore, various models based on the Schrödinger equation have been developed in the literature [2-9]. The stochastic NLSEs which contain the stochastic term with multiplicative noise have also a substantial role in engineering and scientific fields. So, stochastic examinations on NLSE have been especially performed utilizing analytic approaches in recent times. Some of these are as follows: Neirameh and Es-lami examined stochastic chiral NLSE in Ref. [10]. Zayed *et al.*, studied the stochastic Sasa-Satsuma equation involving a multiplicative noise [11]. Cakicioglu *et al.* introduced the stochastic dispersive Schrödinger-Hirota equation

with parabolic law nonlinearity [12]. Zayed *et al.*, discussed (2+1)-dimensional NLSE with spatio-temporal dispersion (STD) [13]. Secer gained the stochastic optical soliton of NLSE with Kerr law nonlinearity in Ref. [14]. In Ref. [15], Mohamed *et al.*, generated new optical soliton solutions of the perturbed stochastic NLSE having generalized anti-cubic nonlinearity. Stochastic solitons to Biswas-Arshed equation with multiplicative white noise were earned in Refs. [16, 17], which includes the Wick-type stochastic NLSE with the conformable derivative. [18] presents the stochastic dark solitons to higher-order NLSE. Additionally, optical soliton solutions of various forms of NLSEs have been derived taking into account different conditions in Refs. [3,19-34].

The Biswas-Milovic equation (BME) [30], a generalized version of the NLSE, is of key significance in modeling the wave propagation of soliton transmission over long distances. Moreover, various versions of the BME have been improved and examined in the literature. Some of these are (2+1)-dimensional BME having Kerr, power, and parabolic law nonlinearities [35], (2+1) and (3+1)-dimensional BME [36], BME with STD and parabolic law [37], BME in presence dual-power law nonlinearity and multiplicative white noise via Itô calculus [38], the perturbed BME with Kudryashov's law of refractive index [39], BME having Kerr law, parabolic law, power law, and dual power-law nonlinearities [40], BME in presence quadratic-cubic and parabolic nonlinearities [41], time fractional BME [42, 43], and BME with STD [44].

In this paper, we aim to explore the optical soliton solutions of the stochastic perturbed BME with cubic-quintic-septic law in the presence of STD and chromatic dispersion having multiplicative white noise in Itô sense via subversion of the new extended auxiliary equation method (SAEM246) [45].

Governing model

The adopted stochastic form of perturbed Biswas-Milovic equation with cubic-quintic-septic law having chromatic dispersion and spatio-temporal in the presence of multiplicative white noise in Itô sense is put forth, for the first time, as:

$$\begin{aligned} i(\vartheta^n)_t + \alpha(\vartheta^n)_{xt} + \beta(\vartheta^n)_{xx} + (c_1|\vartheta|^2 + c_2|\vartheta|^4 \\ + c_3|\vartheta|^6)\vartheta^n = i(\lambda(|\vartheta|^2\vartheta^n)_x + \tau(|\vartheta|^2)_x\vartheta^n \\ + \mu|\vartheta|^2(\vartheta^n)_x) - \sigma(\vartheta^n - i\alpha(\vartheta^n)_x)W_t(t), \end{aligned} \quad (1)$$

in which n is BME to NLSE generalization parameter, α and β symbolize the STD and group velocity dispersion coefficients, c_1, c_2, c_3 express cubic, quintic, and septic law of nonlinearities of self-phase modulation, respectively. Furthermore, τ is defined as a self-frequency shift coefficient whereas λ , and μ are the coefficients of self-steepening and nonlinear terms, respectively. Herein, σ stands for the noise strength, and $W(t)$ is the standard Wiener process which is expressed as [46]:

$$W(t) = \int_0^t \Phi(\eta)dW(\eta), \quad \eta < t, \quad (2)$$

where η is defined as the stochastic variable. In particular, stochastic studies on equations modeling optical soliton behavior come into prominence and become widespread in recent years. However, stochastic studies on the BME are scarcely any. Regarding the perturbed BME, there has been no research reflected in the literature. In this aspect, as far as we know, Eq. (1) has not been introduced before in the literature.

The paper is arranged into five sections. Section 2 expresses the mathematical analysis of Eq. (1). We describe SAEM246 and its utilization to gain the soliton solutions of the stochastic model in Sec. 3. Section 4 summarizes the main findings and their significance. In Sec. 5, conclusions are given.

2. Mathematical analysis of Eq. (1)

The following transformation is taken into account for the model in Eq. (1):

$$\begin{aligned} \vartheta(x, t) &= \mathcal{U}(\zeta) e^{i\phi(x, t)}, \\ \phi(x, t) &= -kx + \omega t + \sigma(W(t) - \sigma t) + \varphi_0, \\ \zeta &= x - \nu t, \end{aligned} \quad (3)$$

in which $\vartheta(x, t)$ is the soliton pulse profile, $\mathcal{U}(\zeta)$ states real function which represents amplitude, φ_0 is phase constant, and k, ω, ν are frequency, wave number, and speed, which are non-zero real values. Inserting Eq. (3) into Eq. (1), the imaginary and real parts of the resultant equation are presented in the following structure, respectively:

$$n \left(\sigma\alpha(n-1) \frac{dW(t)}{dt} + ((k\nu - \sigma^2 + \omega)\alpha - 2\beta k)n - \nu \right) \mathcal{U}'\mathcal{U}^{n+1} - ((\lambda + \mu)n + 2\lambda + 2\tau)\mathcal{U}'\mathcal{U}^{n+3} = 0, \quad (4)$$

$$\begin{aligned} \left(\sigma(n-1)(\alpha kn - 1) \frac{d}{dt} W(t) - n((\sigma^2 - \omega)\alpha + k\beta)kn - \sigma^2 + \omega \right) \mathcal{U}^{n+2} + c_2\mathcal{U}^{n+6} \\ + (c_1 - k(\lambda + \mu)n)\mathcal{U}^{n+4} + c_3\mathcal{U}^{n+8} + n\mathcal{U}''\mathcal{U}^{n+1} - n(n-1)(\alpha\nu - \beta)\mathcal{U}^n(\mathcal{U}')^2 = 0, \end{aligned} \quad (5)$$

where $U = \mathcal{U}(\zeta)$ and Eq. (4) presents the following constraints:

$$\nu = \frac{(\alpha\sigma^2 - \alpha\omega + 2\beta k)n}{\alpha kn - 1}, \quad \tau = -\frac{(\lambda + \mu)n + 2\lambda}{2}. \quad (6)$$

Herein, we need to add the following comment regarding Eq. (4). Considering the first term of Eq. (4), we derive the following expression as an algebraic condition:

$$\left(\sigma\alpha(n-1) \frac{dW(t)}{dt} \right) \mathcal{U}'\mathcal{U}^{n+1} = 0. \quad (7)$$

In Eq. (7), \mathcal{U}' and \mathcal{U}^{n+1} cannot be zero because \mathcal{U} is non-zero and has the second-order derivative as a solution function. Since $W(t)$ has a first-order derivative as the Wiener process, it is not possible for $dW(t)/dt$ to be zero. Because

σ is noise strength and α is the model parameter, they are not zero. Therefore, n must be 1. So, it can be simply interpreted that the cubic-quintic-septic form of Eq. (1) is considered as the NLSE form for BME. Moreover, this evaluation is presented for the first time in this study. So, by considering the $n = 1$, Eq. (5) turns into the following form:

$$\begin{aligned} -((\sigma^2 - \omega)\alpha + k\beta)k - \sigma^2 + \omega \mathcal{U}^3 + c_2\mathcal{U}^7 \\ + (c_1 - k(\lambda + \mu))\mathcal{U}^5 + c_3\mathcal{U}^9 + \mathcal{U}''\mathcal{U}^2 = 0. \end{aligned} \quad (8)$$

Balancing the terms $\mathcal{U}''\mathcal{U}^2$, \mathcal{U}^9 in (8) by considering Eqs. (13), (14), the balance constant is determined as $J=1/3$.

So, we obtain the requirement to perform the following transformation:

$$U(\zeta) = \mathcal{V}^{1/3}(\zeta). \tag{9}$$

Rearrangement of Eq. (8) and using Eq. (9) yields the following nonlinear ordinary differential equation:

$$9(c_1 - (\lambda + \mu)k) \mathcal{V}(\eta)^{\frac{8}{3}} + 9c_2 \mathcal{V}^{\frac{10}{3}} + 9(c_3 \mathcal{V}^2 - \beta k^2 + \alpha(\omega - \sigma^2)k + \sigma^2 - \omega) \mathcal{V}^2 + 2(\alpha\nu - \beta)(\mathcal{V}')^2 - 3(\alpha\nu - \beta)\mathcal{V}\mathcal{V}'' = 0. \tag{10}$$

For integrability of Eq. (10), we must set,

$$c_1 = k(\lambda + \mu), \quad c_2 = 0. \tag{11}$$

As a result, the NLODE in Eq. (10) is formed as follows:

$$9(c_3 \mathcal{V}^2 + \alpha(\omega - \sigma^2)k - \beta k^2 + \sigma^2 - \omega) \mathcal{V}^2 + 2(\alpha\nu - \beta)(\mathcal{V}')^2 - 3(\alpha\nu - \beta)\mathcal{V}\mathcal{V}'' = 0. \tag{12}$$

Balancing the terms $\mathcal{V}''\mathcal{V}$, \mathcal{V}^4 in Eq. (12), the balance constant is determined as positive integer.

3. A subversion of the new extended auxiliary equation method (SAEM246)

The SAEM246 [45] is derived from the subversion of well known auxiliary equation [47-49]. The method offers that Eq. (12) has a solution as the following truncated series form:

$$\mathcal{V}(\zeta) = \sum_{j=0}^J \Lambda_j \mathcal{N}^j(\zeta), \quad \Lambda_J \neq 0, \tag{13}$$

in which Λ_j are real constants, J is the positive integer balance constant to be computed considering the Eqs. (12), (13), and (14) which is obtained as 2. The function $\mathcal{N}(\zeta)$ admits:

$$\left(\frac{d\mathcal{N}}{d\zeta}\right)^2 = \rho^2 \mathcal{N}^2(\zeta) [1 - \chi_1 \mathcal{N}^2(\zeta) - \chi_2 \mathcal{N}^4(\zeta)], \tag{14}$$

in which ρ, χ_1, χ_2 are nonzero real values. One of the solutions for Eq. (14) is given with the following structure [45]:

$$\mathcal{N}(\zeta) = \sqrt{\frac{16e^{2\rho\zeta}}{(e^{2\rho\zeta} + 4\chi_1)^2 + 64\chi_2}}. \tag{15}$$

Considering that J is 2, Eq. (13) is rewritten in the following form:

$$\mathcal{V}(\zeta) = \Lambda_0 + \Lambda_1 \mathcal{N}(\zeta) + \Lambda_2 \mathcal{N}^2(\zeta), \quad \Lambda_2 \neq 0. \tag{16}$$

Inserting Eq. (16) and its corresponding derivatives in Eq. (12) and taking the identical various powers of $\mathcal{N}(\zeta)$ as zero, the following system is earned:

$$\begin{aligned} \mathcal{N}^0(\zeta) : & \Lambda_0^2 \Omega_1 = 0, \\ \mathcal{N}^1(\zeta) : & (\rho^2 \Omega_2 - 6c_3 \Lambda_0^2 + 6\Omega_1) \Lambda_0 \Lambda_1 = 0, \\ \mathcal{N}^2(\zeta) : & (\rho^2 \Lambda_1^2 + 12\rho^2 \Lambda_0 \Lambda_2) \Omega_2 - 9c_3 \Lambda_0^2 (\Omega_3 + 4\Lambda_1^2) - 9\Omega_1 \Omega_3 = 0, \\ \mathcal{N}^3(\zeta) : & (15\rho^2 \Omega_2 - 18c_3 \Lambda_0^2 - 18\Omega_1 - 8\rho^2) \Lambda_1 \Lambda_2 - 6(\rho^2 \chi_1 \Omega_2 + 6c_3 \Omega_3) \Lambda_0 \Lambda_1 = 0, \\ \mathcal{N}^4(\zeta) : & 2\rho^2 \Omega_2 (2\Lambda_2^2 - 4\chi_1 \Lambda_1^2 - 9\chi_1 \Lambda_0 \Lambda_2) - 9c_3 (\Lambda_0^2 \Lambda_2^2 + 8\Lambda_0 \Lambda_1^2 \Lambda_2 + \Omega_3^2) - 9\Lambda_2^2 \Omega_1 = 0, \\ \mathcal{N}^5(\zeta) : & 16\rho^2 \chi_1 \Omega_2 \Lambda_1 \Lambda_2 + 9\rho^2 \chi_2 \Omega_2 \Lambda_0 \Lambda_1 + 36c_3 \Lambda_0 \Lambda_1 \Lambda_2^2 + 36c_3 \Omega_3 \Lambda_1 \Lambda_2 = 0, \\ \mathcal{N}^6(\zeta) : & 10\rho^2 \chi_1 \Omega_2 \Lambda_2^2 + 7\rho^2 \chi_2 \Omega_2 \Lambda_1^2 + 24\rho^2 \chi_2 \Omega_2 \Lambda_0 \Lambda_2 + 36c_3 \Lambda_0 \Lambda_2^3 + 54c_3 \Lambda_1^2 \Lambda_2^2 = 0, \\ \mathcal{N}^7(\zeta) : & (25\rho^2 \chi_2 \Omega_2 + 36c_3 \Lambda_2^2) \Lambda_1 \Lambda_2 = 0, \\ \mathcal{N}^8(\zeta) : & (16\rho^2 \chi_2 \Omega_2 + 9c_3 \Lambda_2^2) \Lambda_2^2 = 0, \end{aligned} \tag{17}$$

in which

$$\Omega_1 = \Lambda_0^2 c_3 + \alpha(\omega - \sigma^2)k - \beta k^2 + \sigma^2 - \omega, \quad \Omega_2 = \left(-\frac{\alpha((-\sigma^2 + \omega)\alpha - 2k\beta)}{k\alpha - 1} - \beta \right),$$

and

$$\Omega_3 = (2\Lambda_0 \Lambda_2 + \Lambda_1^2).$$

Solving the system in Eq. (17), we get:

$$\left\{ \begin{array}{l} \omega = \frac{\sigma^2(\Upsilon + 9\alpha^2k^2) + 4\beta\rho^2(\alpha k + 1) + 9k^2\beta(\alpha k - 1)}{\Upsilon} \\ \chi_1 = 0, \Lambda_0 = 0, \Lambda_1 = 0, \Lambda_2 = -\frac{4\sqrt{c_3\Upsilon\beta\chi_2}\rho}{c_3\Upsilon} \end{array} \right\}. \quad (18)$$

where $\Upsilon = 4\alpha^2\rho^2 + 9\alpha^2k^2 - 18\alpha k + 9$ and $c_3\Upsilon\beta\chi_2 > 0$. Unification of Eqs. (3), (9) (15),(16) allows to reach the solution of Eq. (1):

$$\vartheta_1(x, t) = 16^{\frac{1}{3}} \left(\frac{\Lambda_2 e^{2\rho \left(x + \frac{((\omega - \sigma^2)\alpha - 2k\beta)t}{\alpha k - 1} \right)}}{\left(e^{2\rho \left(x + \frac{((\omega - \sigma^2)\alpha - 2k\beta)t}{\alpha k - 1} \right)} + 4\chi_1 \right)^2 + 64\chi_2} \right)^{\frac{1}{3}} e^{i(-kx + \omega t + \psi_0 + \sigma(W(t) - \sigma t))}, \quad (19)$$

where $\omega, \chi_1, \Lambda_2$ are given in Eq. (18).

4. Results and discussion

This section consists of various graphics indicating the effectiveness of SAEM246 in gaining soliton solutions. Figure 1 consists of nine subfigures for the parameters $\alpha = 1, \beta = c_3 = k = 0.5, \rho = 0.2, \chi_2 = 2, \psi_0 = \sigma = 0$: a) expresses the 3D square of the modulus, b) depicts the contour view, c) illustrates the 2D square of the modulus, d) indicates the 3D imaginary component,




FIGURE 1. Various views of $\vartheta_1(x, t)$ for $\alpha = 1, \beta = c_3 = k = 0.5, \rho = 0.2, \chi_2 = 2, \psi_0 = \sigma = 0$.




FIGURE 2. The various projections of $\vartheta_1(x, t)$ under the effect of noise effect.

e) demonstrates the contour view for the imaginary component, f) represents the 2D imaginary component, g) indicates the 3D real component, h) illustrates the contour view for the real component, i) depicts the 2D real component. Figure 1a) and 1c) display the bright solution of $\vartheta_1(x, t)$ in Eq. (19).

Figure 2 is devoted to the examination of the impact of σ . Figures 2a) and 2b) state the depictions of the real and imaginary parts of Eq. (19), respectively, where the effect of σ can be clearly observed. This effect is seen more clearly when the graphs drawn for $\sigma = 4$ are compared with graphs indicated in Figs. 1d)-1g) for $\sigma = 0$. The resulting effect demonstrates itself in the structure of fluctuations. However, in order to understand the impact of σ more clearly, that is, the situation occurring for increasing values of σ , 2D portraits are drawn. In Figs. 2c) and 2d), we indicate 2D projections of $Re(\vartheta_1(x, t))$ and $Im(\vartheta_1(x, t))$ for $\sigma = 0, 1, 2, 3, 4$. In Fig. 2d), firstly, the graph of Eq. (1) for $\sigma = 0$ (there is no noise effect) is depicted (blue line). This depiction basically has the same character as the wave behavior with Fig. 1i). Then, the values of 1, 2, 3, and 4 for σ were given gradually and the $Re(\vartheta_1(x, t))$ graph was hidden each time (solid red to purple lines) in Fig. 2c). From Fig. 2c), oscillation (fluctuation) is observed with a decrease in the amplitude of the

wave depending on the increasing σ values, and this fluctuation decreases as σ approaches to zero. It is possible to observe the effect for other values of σ which are not limited to the $\sigma = 4$ given in the graphic drawings. A similar examination with Fig. 2d) has been carried out for $Im(\vartheta_1(x, t))$. This study also points out a situation similar to the comments made for $Re(\vartheta_1(x, t))$. Therefore, as a general interpretation, it has been observed that for increasing values of σ , it has a fluctuating impact on the soliton in diverse values of t . Investigations on whether this effect causes a constantly increasing fluctuation depending on increasing σ values, and whether the fluctuation continues (or decreases) after a certain σ value, are some of the topics that can be researched for both the model presented in this study and other optical models.

5. Conclusion

In this research paper, the optical solitons of the stochastic perturbed Biswas-Milovic equation with cubic-quintic-septic law in the presence of STD and chromatic dispersion having by multiplicative white noise in Itô sense were scrutinized via SAEM246. We successfully reached the bright soliton.

3D, contours, and 2D depictions were additionally drawn to figure out the behavior of soliton for the generated solution. Moreover, we intended to analyze the noise influence in terms of Itô sense on soliton solution and presented the depictions of the resultants in detail. In this research paper, it has been observed that the effect of σ is indicated as a fluctuation on the soliton, this fluctuation increases depending on the increasing σ values or decreases when it approaches the value

of $t \neq 0$. For both Eq. (1) and other equations, whether the σ effect will continue as a continuous fluctuation depending on the increasing σ value, whether there is a critical value where the fluctuation stops, and the analysis of the fractional order form of the Eq. (1) are open topics that can be researched in the future. Because of introducing this stochastic problem for the first time in this article, it reveals that the work will conduce to some researchers working in this field.

1. A. Biswas, D. Milovic, and M. Edwards, Mathematical theory of dispersion-managed optical solitons, (Springer Science & Business Media, 2010).
2. E. M. E. Zayed *et al.*, Optical solitons and conservation laws associated with kudryashov's sextic power-law nonlinearity of refractive index, *Ukrainian Journal of Physical Optics* **22** (2021) 38. <https://doi.org/10.3116/16091833/22/1/38/2021>.
3. Y. Yildirim *et al.*, Highly dispersive optical soliton perturbation with kudryashov's sextic-power law of nonlinear refractive index., *Ukrainian Journal of Physical Optics* **23** (2022) 24, <https://doi.org/10.3116/16091833/23/1/24/2022>.
4. A. R. Adem *et al.*, Stationary optical solitons with nonlinear chromatic dispersion for lakshmanan-porsezian-daniel model having kerr law of nonlinear refractive index, *Ukr. J. Phys. Opt.* **22** (2021) 83, <https://doi.org/10.3116/16091833/22/2/83/2021>.
5. A. Biswas, J. Edoki, P. Guggilla, S. Khan, A. K. Alzahrani, M. R. Belic, Cubic-quartic optical soliton perturbation with lakshmanan-porsezian-daniel model by semi-inverse variational principle, *Ukr. J. Phys. Opt* **22** (2021) 123.
6. A. Al Qarni, A. Bodaqah, A. Mohammed, A. Alshaery, H. Bakodah, A. Biswas, Cubic-quartic optical solitons for lakshmanan-porsezian-daniel equation by the improved adomian decomposition scheme, *Ukr. J. Phys. Opt.* **23** (2022) 228.
7. Y. Yildırım *et al.*, Optical solitons in fibre bragg gratings with third-and fourth-order dispersive reflectivities, *Ukr. J. Phys. Opt.* **22** (2021) 239.
8. Y. Yildırım *et al.*, Cubic-quartic optical solitons having quadratic-cubic nonlinearity by sine-gordon equation approach., *Ukrainian Journal of Physical Optics* **22** (2021) 255. <https://doi.org/10.3116/16091833/22/4/255/2021>.
9. O. González-Gaxiola, A. Biswas, Y. Yildırım, H. M. Alshehri, Highly dispersive optical solitons in birefringent fibres with non) local form of nonlinear refractive index: Laplace-adomian decomposition., *Ukrainian Journal of Physical Optics* **23** (2022) 68, <https://doi.org/10.3116/16091833/23/2/68/2022>.
10. A. Neirameh, M. Eslami, New optical soliton of stochastic chiral nonlinear Schrödinger equation, *Optical and Quantum Electronics* **55** (2023) 444, <https://doi.org/10.1007/s11082-023-04564-8>.
11. E. M. Zayed *et al.*, Optical solitons in the sasa-satsuma model with multiplicative noise via Itô calculus, *Ukr. J. Phys. Opt.* **23** (2022) 9, <https://doi.org/10.3116/16091833/23/1/9/2022>.
12. H. Cakicioglu, M. Ozisik, A. Secer, M. Bayram, Stochastic dispersive Schrödinger-hirota equation having parabolic law nonlinearity with multiplicative white noise via ito calculus, *Optik* **279** (2023) 170776, <https://doi.org/10.1016/j.ijleo.2023.170776>.
13. E. M. Zayed, M. E. Alngar, R. M. Shohib, K. A. Gepreel, T. A. Nofal, Optical solitons with (2+1)- dimensional nonlinear Schrödinger equation having spatio-temporal dispersion and multiplicative white noise via ito calculus, *Optik* **261** (2022) 169204. <https://doi.org/10.1016/j.ijleo.2022.169204>.
14. A. Secer, Stochastic optical solitons with multiplicative white noise via Itô calculus, *Optik* **268** (2022) 169831, <https://doi.org/10.1016/j.ijleo.2022.169831>.
15. E. Mohamed, I. El-Kalla, A. Tarabia, A. A. Kader, New optical solitons for perturbed stochastic nonlinear schrödinger equation by functional variable method, *Optical and Quantum Electronics* **55** (2023) 603, <https://doi.org/10.1007/s11082-023-04844-3>.
16. Y. Alhojilan, H. M. Ahmed, W. B. Rabie, Stochastic solitons in birefringent fibers for biswas-arshed equation with multiplicative white noise via Itô calculus by modified extended mapping method, *Symmetry* **15** (2023) 207, <https://doi.org/10.3390/sym15010207>.
17. E. Ulutas, Travelling wave and optical soliton solutions of the wick-type stochastic nlse with conformable derivatives, *Chaos, Solitons & Fractals* **148** (2021) 111052, <https://doi.org/10.1016/j.chaos.2021.111052>.
18. H. Zhong, B. Tian, M. Li, W.-R. Sun, H.-L. Zhen, Stochastic dark solitons for a higher-order nonlinear Schrödinger equation in the optical fiber, *Journal of Modern Optics* **60** (2013) 1644, <https://doi.org/10.1080/09500340.2013.852713>.
19. X. Liu, W. Liu, H. Triki, Q. Zhou, A. Biswas, Periodic attenuating oscillation between soliton interactions for higher-order variable coefficient nonlinear Schrödinger equation, *Nonlinear Dynamics* **96** (2019) 801, <https://doi.org/10.1007/s11071-019-04822-z>.
20. Z.-y. Zhang, Y.-x. Li, Z.-h. Liu, X.-j. Miao, New exact solutions to the perturbed nonlinear Schrödinger's equation with kerr law nonlinearity via modified trigonometric function series method,

- Communications in Nonlinear Science and Numerical Simulation* **16** (2011) 3097, <https://doi.org/10.1016/j.cnsns.2010.12.010>.
21. M. Ozisik *et al.*, Investigation of optical soliton solutions of higher-order nonlinear Schrödinger equation having kudryashov nonlinear refractive index, *Optik* (2023) 170548, <https://doi.org/10.1016/j.ijleo.2023.170548>.
 22. N. Ozdemir, Optical solitons for radhakrishnan-kundulakshmanan equation in the presence of perturbation term and having kerr law, *Optik* **271** (2022) 170127, <https://doi.org/10.1016/j.ijleo.2022.170127>.
 23. N. Ozdemir, A. Secer, M. Ozisik, M. Bayram, Perturbation of dispersive optical solitons with Schrödinger Hirota equation with kerr law and spatio-temporal dispersion, *Optik* **265** (2022) 169545, <https://doi.org/10.1016/j.ijleo.2022.169545>.
 24. N. A. Kudryashov, Hamiltonians of the generalized nonlinear Schrödinger equations, *Mathematics* **11** (2023) 2304, <https://doi.org/10.3390/math11102304>.
 25. N. A. Kudryashov, Embedded solitons of the generalized nonlinear Schrödinger equation with high dispersion, *Regular and Chaotic Dynamics* **27** (2022) 680, <https://doi.org/10.1134/S1560354722060065>.
 26. H. Kacicioglu, M. Ozisik, A. Secer, M. Bayram, Optical soliton solutions of Schrödinger-Hirota equation with parabolic law nonlinearity via generalized kudryashov algorithm, *Optical and Quantum Electronics* **55** (2023) 407, <https://doi.org/10.1007/s11082-023-04634-x>.
 27. I. Onder, A. Secer, M. Bayram, Optical soliton solutions of time-fractional coupled nonlinear Schrödinger system via kudryashov-based methods, *Optik* **272** (2023) 170362, <https://doi.org/10.1016/j.ijleo.2022.170362>.
 28. N. A. Kudryashov, Solitary waves of the generalized radhakrishnan-kundulakshmanan equation with four powers of nonlinearity, *Physics Letters A* **448** (2022) 128327, <https://doi.org/10.1016/j.physleta.2022.128327>.
 29. E. M. Zayed, R. M. Shohib, K. A. Gepreel, M. M. El-Horbaty, and M. E. Alngar, Cubic-quartic optical soliton perturbation biswas-milovic equation with kudryashov's law of refractive index using two integration methods, *Optik* **239** (2021) 166871, <https://doi.org/10.1016/j.ijleo.2021.166871>.
 30. A. Biswas, D. Milovic, Bright and dark solitons of the generalized nonlinear Schrödinger's equation, *Communications in Nonlinear Science and Numerical Simulation* **15** (2010) 1473, <https://doi.org/10.1016/j.cnsns.2009.06.017>.
 31. E. C. Aslan, F. Tchier, and M. Inc, On optical solitons of the Schrödinger-Hirota equation with power law nonlinearity in optical fibers, *Superlattices and Microstructures* **105** (2017) 48, <https://doi.org/10.1016/j.spmi.2017.03.014>.
 32. A. Biswas, M. Ekici, A. Sonmezoglu, and M. R. Belic, Highly dispersive optical solitons with cubic-quintic-septic law by f-expansion, *Optik* **182** (2019) 897, <https://doi.org/10.1016/j.ijleo.2019.01.058>.
 33. A. A. Magazev, M. N. Boldyreva, Schrödinger equations in electromagnetic fields: Symmetries and noncommutative integration, *Symmetry* **13** (2021) 1527, <https://doi.org/10.3390/sym13081527>.
 34. H. Abdelwahed, E. El-Shewy, R. Sabry, M. A. Abdelrahman, Characteristics of stochastic langmuir wave structures in presence of Itô sense, *Results in Physics* **37** (2022) 105435, <https://doi.org/10.1016/j.rinp.2022.105435>.
 35. S. Altun, M. Ozisik, A. Secer, M. Bayram, Optical solitons for biswas-milovic equation using the new kudryashov's scheme, *Optik* **270** (2022) 170045, <https://doi.org/10.1016/j.ijleo.2022.170045>.
 36. M. Ozisik, Novel (2+ 1) and (3+ 1) forms of the biswas-milovic equation and optical soliton solutions via two efficient techniques, *Optik* **269** (2022) 169798, <https://doi.org/10.1016/j.ijleo.2022.169798>.
 37. P. Albayrak, Optical solitons of biswas-milovic model having spatio-temporal dispersion and parabolic law via a couple of kudryashov's schemes, *Optik* **279** (2023) 170761, <https://doi.org/10.1016/j.ijleo.2023.170761>.
 38. E. M. Zayed, R. M. Shohib, M. E. Alngar, Dispersive optical solitons with biswas-milovic equation having dual-power law nonlinearity and multiplicative white noise via Itô calculus, *Optik* **270** (2022) 169951, <https://doi.org/10.1016/j.ijleo.2022.169951>.
 39. L. Akinyemi, M. Mirzazadeh, S. Amin Badri, K. Hosseini, Dynamical solitons for the perturbed biswas-milovic equation with kudryashov's law of refractive index using the first integral method, *Journal of Modern Optics* **69** (2022) 172, <https://doi.org/10.1080/09500340.2021.2012286>.
 40. N. Raza, M. Abdullah, A. R. Butt, Analytical soliton solutions of biswas-milovic equation in kerr and non-kerr law media, *Optik* **157** (2018) 993, <https://doi.org/10.1016/j.ijleo.2017.11.043>.
 41. B. Ghanbari *et al.*, Families of exact solutions of biswas-milovic equation by an exponential rational function method, *Tbilisi Mathematical Journal* **13** (2020) 65, <https://doi.org/10.32513/tbilisi/1593223219>.
 42. B. Kour, S. Kumar, Time fractional biswas-milovic equation: Group analysis, soliton solutions, conservation laws and residual power series solution, *Optik* **183** (2019) 1085, <https://doi.org/10.1016/j.ijleo.2019.02.099>.
 43. P. Sunthrayuth, M. Naeem, N. A. Shah, R. Shah, J. D. Chung, On the solution of fractional biswas-milovic model via analytical method, *Symmetry* **15** (2023) 210, <https://doi.org/10.3390/sym15010210>.
 44. N. Raza, A. Javid, Optical dark and singular solitons to the biswas-milovic equation in nonlinear optics with spatio-temporal dispersion, *Optik* **158** (2018) 1049, <https://doi.org/10.1016/j.ijleo.2017.12.186>.
 45. M. Ozisik, A. Secer, M. Bayram, H. Aydin, An encyclopedia of kudryashov's integrability approaches applicable to optoelectronic devices, *Optik* **265** (2022) 169499, <https://doi.org/10.1016/j.ijleo.2022.169499>.

46. W. W. Mohammed, H. Ahmad, H. Boulares, F. Khelifi, M. El-Morshedy, Exact solutions of hirota- maccari system forced by multiplicative noise in the Itô sense, *Journal of Low Frequency Noise, Vibration and Active Control* **41** (2022) 74, <https://doi.org/10.1177/14613484211028100>.
47. E. Yomba, A generalized auxiliary equation method and its application to nonlinear klein-gordon and generalized nonlinear camassa-holm equations, *Physics Letters A* **372** (2008) 1048, <https://doi.org/10.1016/j.physleta.2007.09.003>.
48. Sirendaoreji, A new auxiliary equation and exact traveling wave solutions of nonlinear equations, *Physics Letters A* **356** (2006) 124, <https://doi.org/10.1016/j.physleta.2006.03.034>.
49. E. M. Zayed, A further improved (g'/g)-expansion method and the extended tanh-method for finding exact solutions of nonlinear pdes, *Wseas Transactions on Mathematics* **10** (2011) 56.