

ERRATUM

The n -particle fractional parentage coefficients for anticommuting creation operators, C. Quesne, Rev. Mex. Fís. 19 (1970) 109.

Formulae (13), (14) and (15) should read respectively:

$$\begin{aligned}
 & |m'_1 \dots m'_{n'} m_1 \dots m_n |_{1 \dots n+n'} \\
 &= \left\{ \binom{n+n'}{n} \right\}^{-\frac{1}{2}} \sum_p (-1)^p |m'_1 \dots m'_{n'} |_{p(1) \dots p(n')} \times \\
 & \times |m_1 \dots m_n |_{p(n'+1) \dots p(n+n')} , \tag{13}
 \end{aligned}$$

$$\begin{aligned}
 & \left[P_{n' \gamma' J'}^+ P_{n \gamma J}^+ \right]_{J'' M''} |0\rangle = \left\{ \binom{n+n'}{n} \right\}^{-\frac{1}{2}} \sum_p (-1)^p \\
 & \times \psi \left[\begin{matrix} j^{n'} \\ p(1) \dots p(n') \end{matrix} (\gamma' J') j^n \begin{matrix} j^n \\ p(n'+1) \dots p(n+n') \end{matrix} (\gamma J) J'' M'' \right] \tag{14}
 \end{aligned}$$

$$\langle j^{n'} \gamma' J' ; j^n \gamma J | j^{n+n'} \gamma'' J'' \rangle = \left\{ \binom{n+n'}{n} \right\}^{-\frac{1}{2}} \frac{(-1)^{n(n'+1)}}{\sqrt{2J''+1}}$$

$$\times \langle n'' \gamma'' J'' || P_{n \gamma J}^+ || n' \gamma' J' \rangle . \tag{15}$$